



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON RESTRICTED SYSTEMS OF HIGHER INDETERMINATE EQUATIONS *

BY

E. T. BELL

By two examples we shall illustrate a means for deriving arithmetical properties of certain restricted forms. So little being known concerning the arithmetic of forms of higher degree, it is reasonable to develop what is offered by analysis. The method exemplified is of wide scope and easy applicability, and may possibly, in addition to giving new properties of integers, yield also, on starting from the appropriate associated forms of the second degree, information regarding current problems. It is a natural extension to forms of higher degree of the classic applications of elliptic functions to special quadratic forms. The method being new, we shall give the analysis for the first example in some detail.

1. We take first a single restricted form

$$(1) \quad uy^s + vx^r - x^r y^s$$

in which u, v, x, y are variable integers; u, v are odd, x, y even, and r, s are constant odd positive integers. For $\alpha \geq 1$ and m odd and positive, we consider all the representations (u, v, x, y) of $2^\alpha m$ in (1); or, what is the same, the totality of solutions (u, v, x, y) of

$$(2) \quad 2^\alpha m = uy^s + vx^r - x^r y^s,$$

where α, m, r, s are constants, m, u, v, r, s are odd, x, y even, and $\alpha \geq 1$. From all the (u, v, x, y) we select two classes C_1, C_2 defined by

$$\begin{aligned} (C_1): \quad & x \equiv 0, \quad y > 0, \quad u > x^r, \quad v < y^s, \\ (C_2): \quad & x > 0, \quad y \equiv 0, \quad u < x^r, \quad v > y^s; \end{aligned}$$

and denote by $N_i(2^\alpha m)$ the number of (u, v, x, y) belonging to C_i . It will be shown that the difference $N_1(2^\alpha m) - N_2(2^\alpha m)$ is a function of α, r, s and the divisors of m .

2. It is convenient to distinguish four cases:

$$\begin{aligned} (A_1): \quad & \alpha \equiv 0 \pmod{r} \quad \text{and} \quad \alpha \equiv 0 \pmod{s}, \\ (A_2): \quad & \alpha \equiv 0 \pmod{r} \quad \text{and} \quad \alpha \not\equiv 0 \pmod{s}, \\ (A_3): \quad & \alpha \not\equiv 0 \pmod{r} \quad \text{and} \quad \alpha \equiv 0 \pmod{s}, \\ (A_4): \quad & \alpha \not\equiv 0 \pmod{r} \quad \text{and} \quad \alpha \not\equiv 0 \pmod{s}. \end{aligned}$$

* Presented to the Society, October, 1920.

Let $\lambda_j(n)$ denote the sum of all those divisors of the positive integer n which are j th powers. Then, corresponding to the four cases we have:

$$\begin{aligned}(A'_1): \quad & N_1(2^\alpha m) - N_2(2^\alpha m) = 2^{\alpha-1} [\lambda_s(m) - \lambda_r(m)], \\(A'_2): \quad & N_1(2^\alpha m) - N_2(2^\alpha m) = -2^{\alpha-1} \lambda_r(m), \\(A'_3): \quad & N_1(2^\alpha m) - N_2(2^\alpha m) = 2^{\alpha-1} \lambda_s(m), \\(A'_4): \quad & N_1(2^\alpha m) - N_2(2^\alpha m) = 0,\end{aligned}$$

all of which may be put in a single statement. To prove these results we transform (1) into an associated quadratic form, simultaneously replacing the conditions (C_1) , (C_2) by equivalent restrictions on the new variables.

3. The (u, v, x, y) being as in § 1, we see that d' , δ' , d'' , δ'' defined by

$$(3) \quad u = d', \quad v = \delta', \quad u - x^r = d'', \quad y^s - v = \delta'',$$

are odd positive integers provided that (u, v, x, y) belongs to C_1 . Let (u, v, x, y) belong to C_1 . Then, since

$$uy^s + vx^r - x^r y^s = uv + (u - x^r)(y^s - v),$$

we may replace (2) by

$$(4) \quad 2^\alpha m = d' \delta' + d'' \delta'',$$

in which d' , δ' , d'' , δ'' are restricted to be odd positive integers subject to the conditions

$$(5) \quad d' - d'' = x^r, \quad \delta' + \delta'' = y^s, \quad x \geq 0,$$

and, necessarily, $y > 0$. It is easily seen that all those solutions of (2) belonging to C_1 are identical with all those of (4) which satisfy also (5). That is, (2) and (C_1) define the same (u, v, x, y) as do (4) and (5). Hence $N_1(2^\alpha m)$ is equal to the number of solutions $(d', \delta', d'', \delta'')$ of (4) such that $d' - d''$ is the r th power of an integer ≥ 0 , and $\delta' + \delta''$ the s th power of an integer > 0 , the d' , δ' , d'' , δ'' being odd positive integers as defined.

Similarly, starting from

$$u = d', \quad v = \delta', \quad x^r - u = d'', \quad v - y^s = \delta'',$$

where d' , δ' , d'' , δ'' are odd positive integers, and (u, v, x, y) belongs to C_2 , we see that all the solutions of (2) belonging to C_2 are identical with all those of (4) which satisfy also

$$d' + d'' = x^r, \quad \delta' - \delta'' = y^s, \quad x > 0, \quad y \geq 0;$$

that is, $N_2(2^\alpha m)$ is equal to the number of solutions $(d', \delta', d'', \delta'')$ of (4) which are such that $d' + d''$ is the r th power of an integer > 0 , and $\delta' - \delta''$ the s th power of an integer ≥ 0 .

4. To evaluate $N_i(2^\alpha m)$ from this reduction to an associated quadratic form, we remark that if n is the k th power (k an integer) of an integer, $-n$ is also the k th power of an integer when and only when k is odd. Hence if $\varphi_{2t-1}(x)$ denotes 1 or 0 according as x is or is not the $(2t-1)$ th power of an integer, $\varphi_{2t-1}(x)$ is an even function of x , viz.,

$$\varphi_{2t-1}(x) = \varphi_{2t-1}(-x).$$

Again, if x, y are relatively prime,

$$\varphi_{2t-1}(xy) = \varphi_{2t-1}(x) \varphi_{2t-1}(y);$$

and if x, y, z, \dots are integers, r, s, t, \dots odd positive integers, the value of

$$\varphi_{r,s,t,\dots}(x,y,z,\dots), \equiv \varphi_r(x) \varphi_s(y) \varphi_t(z) \dots$$

is unity when and only when simultaneously x is an r th power, y is an s th power, z is a t th power, \dots , and in every other case the function vanishes. That is, this function is even in each of its variables x, y, z, \dots , and vanishes with each of $\varphi_r, \varphi_s, \varphi_t, \dots$.

5. Using the values of $N_i(2^\alpha m)$ deduced in § 3 from the associate (4), we have now

$$\begin{aligned} N_1(2^\alpha m) &= \sum \varphi_{r,s}(d' - d'', \delta' + \delta''), \\ N_2(2^\alpha m) &= \sum \varphi_{r,s}(d' + d'', \delta' - \delta''), \end{aligned}$$

the \sum extending to all the $d', \delta', d'', \delta''$ given by (4) as defined in § 3. Let $f(x, y)$ denote any function which is even in each of its variables x, y . Then, the \sum on the left being as just stated, that on the right extending to all positive divisors d of m , there is the well-known result due to Liouville:

$$\begin{aligned} \sum [f(d' - d'', \delta' + \delta'') - f(d' + d'', \delta' - \delta'')] \\ = 2^{\alpha-1} \sum d [f(0, 2^\alpha d) - f(2^\alpha d, 0)]. \end{aligned}$$

Hence, substituting $\varphi_{r,s}(x, y)$ for $f(x, y)$, we have

$$(6) \quad N_1(2^\alpha m) - N_2(2^\alpha m) = 2^{\alpha-1} \sum d [\varphi_{r,s}(0, 2^\alpha d) - \varphi_{r,s}(2^\alpha d, 0)];$$

and since d is odd,

$$\varphi_r(2^\alpha d) = \varphi_r(2^\alpha) \varphi_r(d), \quad \varphi_s(2^\alpha d) = \varphi_s(2^\alpha) \varphi_s(d),$$

so that (6) is equivalent to $(A'_1), \dots, (A'_4)$ of § 2.

6. In deriving such results the order of the steps is the reverse of that just completed. It will be sufficiently evident from the following example. The function $f(x, y)$ is as in § 5, and all the letters, unless otherwise noted, represent positive integers. The origin of the fundamental f -identity is indicated in § 8.

Consider the pair of equations

$$(7) \quad \begin{cases} 2n = 2n_1 + 2n_2, & n_1 = t_1 \tau_1, & n_2 = t_2 \tau_2, \\ 2n = m'_1 + m'_2, & m'_1 = t'_1 \tau'_1, & m'_2 = t'_2 \tau'_2, \end{cases}$$

in which the t, τ are pairs of conjugate divisors, as indicated, and the t 's are divisors whose conjugates are odd; n_1, n_2 are odd or even, m'_1, m'_2 are odd. Hence $t'_1, \tau'_1, t'_2, \tau'_2$ are odd, as also are τ_1, τ_2 ; and $t_i (i = 1, 2)$ is odd or even according as n_i is odd or even, that is, t_i and n_i are of the same parity. Then, the \sum on the left referring to all t, τ, m'_2 which satisfy (7) for n given, that on the right to all pairs of conjugate divisors t, τ of $n (= t\tau)$, we have

$$\begin{aligned} & \sum [(-1)^{t_2} \{f(2t_2 + \tau_1, \tau_2) - f(2t_2 - \tau_1, \tau_2)\} \\ & \quad - (-1)^{t_1} \{f(\tau_2, 2t_2 + \tau_1) - f(\tau_2, 2t_2 - \tau_1)\}] \\ & \quad + 2 \sum (-1 | \tau'_1) \xi(m'_2) f(t'_1, \tau'_1) \\ & \quad = \sum_{h=1}^t \{f(2h-1, \tau) - (-1)^n f(\tau, 2h-1)\}, \end{aligned}$$

where $(-1 | \tau'_1)$ is Jacobi's extension of Legendre's symbol, and $\xi(m'_2)$ one fourth the number of representations of m'_2 as a sum of two squares. This identity indicates, on replacing $f(x, y)$ by $\varphi_{r,s}(x, y)$, the substitutions for transforming (7) into four higher forms, also what restrictions are to be imposed on the latter.

7. For this purpose we consider the first partial sum on the left (after replacing f by φ),

$$\sum (-1)^{t_2} \varphi_{r,s}(2t_2 + \tau_1, \tau_2).$$

Here the φ -function = 1 when and only when integers x, y exist such that

$$2t_2 + \tau_1 = x^r, \quad \tau_2 = y^s.$$

In every other case the value is zero. Hence we replace the first of the equations (7) by

$$(8) \quad 2n = 2vx^r + 2uy^s - 4uv,$$

the substitution converting it into (8) being

$$t_2 = u, \quad \tau_1 = x^r - 2u, \quad t_1 = v, \quad \tau_2 = y^s.$$

By translating the conditions upon the t, τ into terms of the new variables, we get the restrictions to which (8) is subjected:

$$(C_1): \quad x, y \text{ odd} > 0; u, v \text{ odd or even}, > 0; x^r > 2u.$$

Further subdividing C_1 according as u is even or odd, we have two subclasses,

$$(C_{10}): \quad x, y \text{ odd}, > 0; u \text{ even}, > 0; v \text{ odd or even}, > 0; x^r > 2u;$$

$$(C_{11}): \quad x, y \text{ odd}, > 0; u \text{ odd}, > 0; v \text{ odd or even}, > 0; x^r > 2u.$$

Let $N_{ij}(2n)$ denote the number of solutions of (8) which belong to C_{ij} . Then, as in the first example, we have

$$\sum (-1)^{t_2} \varphi_{r,s}(2t_2 + \tau_1, \tau_2) = N_{10}(2n) - N_{11}(2n).$$

The third partial sum also refers to (8), the substitution being

$$t_2 = v, \quad \tau_1 = y^s - 2v, \quad t_1 = u, \quad \tau_2 = x^r,$$

and the corresponding classes

$$(C_{30}): \quad x, y \text{ odd, } > 0; u \text{ even, } > 0; v \text{ odd or even, } > 0; y^s > 2v;$$

$$(C_{31}): \quad x, y \text{ odd, } > 0; u \text{ odd, } > 0; v \text{ odd or even, } > 0; y^s > 2v,$$

$$\text{whence } \sum (-1)^{t_1} \varphi_{r,s}(\tau_2, 2t_2 + \tau_1) = N_{30}(2n) - N_{31}(2n).$$

The second partial sum refers to the equation

$$(9) \quad 2n = -2v_1 x_1^r + 2u_1 y_1^s + 4u_1 v_1,$$

the substitution and the classes being, with a notation similar to that for the preceding cases,

$$t_2 = u_1, \quad \tau_1 = 2u_1 - x_1^r, \quad t_1 = v_1, \quad \tau_2 = y_1^s;$$

$$(C_{20}): \quad x_1, y_1 \text{ odd, } x_1 \geq 0, y_1 > 0; u_1 \text{ even, } > 0; v_1 \text{ odd or even, } > 0; x_1^r < 2u_1;$$

$$(C_{21}): \quad x_1, y_1 \text{ odd, } x_1 \geq 0, y_1 < 0; u_1 \text{ odd, } > 0; v_1 \text{ odd or even, } > 0; x_1^r < 2u_1;$$

whence

$$\sum (-1)^{t_2} \varphi_{r,s}(2t_2 - \tau_1, \tau_2) = N_{20}(2n) - N_{21}(2n).$$

The fourth partial sum refers to

$$(10) \quad 2n = 2v_2 x_2^r - 2u_2 y_2^s + 4u_2 v_2,$$

the substitution and classes being

$$t_2 = v_2, \quad \tau_1 = 2v_2 - y_2^s, \quad t_1 = u_2, \quad \tau_2 = x_2^r;$$

$$(C_{40}): \quad x_2, y_2 \text{ odd, } x_2 > 0, y_2 \geq 0; u_2 \text{ even, } > 0; v_2 \text{ odd or even, } > 0; y_2^s < 2v_2;$$

$$(C_{41}): \quad x_2, y_2 \text{ odd, } x_2 > 0, y_2 \geq 0; u_2 \text{ odd, } > 0; v_2 \text{ odd or even, } > 0; y_2^s < 2v_2;$$

whence

$$\sum (-1)^{t_1} \varphi_{r,s}(\tau_2, 2t_2 - \tau_1) = N_{40}(2n) - N_{41}(2n).$$

The fifth partial sum refers to the equation

$$(11) \quad 2n = x_3^r y_3^s + u_3^2 + v_3^2,$$

for which the substitution and classes are

$$t'_1 = x_3^r, \quad \tau'_1 = y_3^s, \quad m'_2 = u_3^2 + v_3^2;$$

$$(C_{50}): \quad x_3, y_3 \text{ odd, } > 0; u_3, v_3 \text{ of opposite parities, } \not\equiv 0; y_3 \equiv 1 \pmod{4};$$

$$(C_{51}): \quad x_3, y_3 \text{ odd, } > 0; u_3, v_3 \text{ of opposite parities, } \not\equiv 0; y_3 \equiv -1 \pmod{4};$$

and we have

$$2 \sum (-1)^{|\tau'_1|} \xi(m'_2) \varphi_{r,s}(t'_1, \tau'_1) = \frac{1}{2} [N_{50}(2n) - N_{51}(2n)].$$

To evaluate the right side of the identity we denote by $[x]$ the greatest integer which does not exceed x , and by t_j a positive divisor of n whose conjugate, τ_j , is a j th power. Then, the \sum on the right referring to all t_r, t_s , we have at once

$$\begin{aligned} \sum_{h=1}^t [\varphi_{r,s}(2h-1, \tau) - (-1)^n \varphi_{r,s}(\tau, 2h-1)] \\ = \sum \{[\sqrt[r]{2t_s} - 1] - (-1)^n [\sqrt[s]{2t_r} - 1]\}. \end{aligned}$$

Combining all these results, we get for the four equations (8), (9), (10), (11) the following syzygy between the number of solutions belonging to the classes C_{ij} :

$$\begin{aligned} 2\{N_{10}(2n) - N_{11}(2n) - N_{20}(2n) + N_{21}(2n) - N_{30}(2n) + N_{31}(2n) \\ + N_{40}(2n) - N_{41}(2n)\} + N_{50}(2n) - N_{51}(2n) \\ = 2 \sum \{[\sqrt[r]{2t_s} - 1] - (-1)^n [\sqrt[s]{2t_r} - 1]\}; \end{aligned}$$

and this may be regarded as a relation between the numbers of simultaneous representations of $2n$ in the system of forms

$$\begin{aligned} 2vx^r + 2uy^s - 4uv; & \quad -2v_1x_1^r + 2u_1y_1^s + 4u_1v_1; \\ 2v_2x_2^r - 2u_2y_2^s + 4u_2v_2; & \quad x_3^s y_3^r + u_3^2 + v_3^2, \end{aligned}$$

subject to the given restrictions. In both examples the number of representations in each class is obviously, from the associate, finite.

8. The identity in § 6 is one paraphrase of the equation of three terms in elliptic functions; Liouville's in § 5 may be derived from the same source, as shown in papers presented to the Society in 1918. By the methods of those papers combined with the present, similar results for systems of any number of forms in any number of indeterminates may be found.

UNIVERSITY OF WASHINGTON,
SEATTLE, WASH.